

HEEGNER DIVISORS IN THE MODULI SPACE OF GENUS THREE CURVES

MICHELA ARTEBANI

ABSTRACT. S. Kondō used periods of $K3$ surfaces to prove that the moduli space of genus three curves is birational to an arithmetic quotient of a complex 6-ball. In this paper we study Heegner divisors in the ball quotient, given by arithmetically defined hyperplane sections of the ball. We show that the corresponding loci of genus three curves are given by: hyperelliptic curves, singular plane quartics and plane quartics admitting certain rational “splitting curves”.

CONTENTS

Introduction	1
1. Notation and Preliminaries	2
2. The model by Kondō	4
3. Heegner divisors and embeddings	8
4. Heegner divisors and genus 3 curves	10
5. Examples	15
Appendix A. Lattices	20
References	22

INTRODUCTION

The degree four cyclic cover of the projective plane branched along a smooth plane quartic is a $K3$ surface endowed with an automorphism group $G \cong \mathbb{Z}_4$. This simple geometric construction relates the moduli space $\mathcal{M}_3$ of genus three curves to a moduli space $\mathcal{M}$ of polarized $K3$ surfaces. A generator σ for G can be chosen such that the period point of the polarized $K3$ surface (X, σ) belongs to the i -eigenspace $W_{\mathbb{C}}$ of σ^* in $H^2(X, \mathbb{C})$. This implies that the period domain is a six dimensional complex ball $B \subset \mathbb{P}(W_{\mathbb{C}})$ and the moduli space $\mathcal{M}$ is obtained by taking the quotient of B by the action of an arithmetic group Γ . The correspondence between genus three curves and polarized $K3$ surfaces is thus described by a period map

$$\mathcal{P} : \mathcal{M}_3 \setminus \mathcal{M}_3^h \longrightarrow \mathcal{M} \cong B/\Gamma,$$

where $\mathcal{M}_3^h$ denotes the hyperelliptic locus. In fact, in [7] S. Kondō proves that this map is birational and induces an isomorphism:

$$\mathcal{M}_3 \setminus \mathcal{M}_3^h \cong \mathcal{M} \setminus (\mathcal{D}_n \cup \mathcal{D}_h),$$

Date: 2nd February 2008.

Key words and phrases. genus three curves, splitting curves, $K3$ surfaces, Heegner divisors.

This work is partially supported by: PRIN 2003: Spazi di moduli e teoria di Lie; GNSAGA.

where $\mathcal{D}_n, \mathcal{D}_h$ are two irreducible divisors, called *mirrors*.

The $\mathbb{Q}$ -vector space $H^2(X, \mathbb{Q})$ and the action of σ give naturally a vector space W over $k = \mathbb{Q}[i]$ such that $W_{\mathbb{C}} = W \otimes_k \mathbb{C}$. An interesting class of divisors in $\mathcal{M}$ is then given by quotients of hyperplane sections of B defined over k . These divisors are again arithmetic quotients of a (five dimensional) complex ball and are called *Heegner divisors*. This kind of divisors has been introduced and studied in a more general setting in [8].

The first examples of Heegner divisors are the two mirrors $\mathcal{D}_n, \mathcal{D}_h$. In [7] it is proved that the generic points in the mirrors correspond, via the period map $\mathcal{P}$, to a plane quartic with a node and to a smooth hyperelliptic genus three curve respectively. The aim of this paper is to describe all Heegner divisors in terms of genus three curves. We prove indeed that any Heegner divisor which is not a mirror can be interpreted as the locus of plane quartics admitting a *splitting curve* i.e. an irreducible plane curve whose inverse image by the degree four cyclic cover is the union of four distinct curves on the $K3$ surface. More precisely, we show that the splitting curve can be chosen to be rational. The simplest example of Heegner divisor of this kind is the divisor of plane quartics with a hyperflex line.

The paper is organized as follows.

In the first section we introduce some basic notations for genus three curves, Del Pezzo surfaces and $K3$ surfaces.

The geometric construction by Kondō and its main theorem are recalled in the second section.

In the third section we define Heegner divisors and show how they are connected to the problem of embedding a rank two lattice in a non-unimodular lattice. In fact, we introduce two natural invariants associated to such embeddings: the type $n \in \mathbb{Z}$, $n > 0$ and the index $m \in \{1, 2\}$. We also provide an existence result for Heegner divisors with fixed type and index.

In the following section Heegner divisors are interpreted in terms of genus three curves. The main theorem states that the generic point in a Heegner divisor of type $n > 1$ corresponds to a smooth plane quartic having a splitting rational curve with smooth preimages in the Del Pezzo surface. Moreover, we give the minimal degree of such a splitting curve as a function of n and m (it turns out that it is even if $m = 1$ and odd if $m = 2$).

Some examples of Heegner divisors are provided in the last section for splitting curves of degree 1, 2 and 3.

The Appendix recalls some results of lattice theory which will be useful in the proofs.

Acknowledgements. I would like to thank Professor B. van Geemen for introducing me to this subject and for several valuable suggestions. I also wish to thank Dr. A. Laface for many interesting discussions. Part of this work was done during a visit to the Queen's University (Kingston, Ontario), I thank the Mathematics Department for hosting me.

1. NOTATION AND PRELIMINARIES

In this section we introduce the three geometric objects of main interest in this paper and fix the relative notations: genus three curves, Del Pezzo surfaces and $K3$

surfaces. Our main references are [9], [4] and [2].

Let $\mathcal{M}_3$ be the coarse moduli variety parametrizing isomorphism classes of smooth *genus three curves*. We denote with $\mathcal{M}_3^h$ the hyperelliptic divisor in $\mathcal{M}_3$.

A *Del Pezzo surface* is a smooth surface with ample anti-canonical bundle. These surfaces can be equivalently defined as the blowing up of $\mathbb{P}^2$ in a set of $m \leq 8$ distinct points “in general position”. We are interested in Del Pezzo surfaces of degree two i.e. the blowing up of $\mathbb{P}^2$ in seven points $p_1, \dots, p_7$. In this case, a natural basis for the Picard lattice of the surface S is given by

$$Pic(S) = \langle e_0, e_1, \dots, e_7 \rangle,$$

where e_0 is the pull-back of the hyperplane bundle of $\mathbb{P}^2$ and e_i is the exceptional divisor corresponding to p_i , $i = 1, \dots, 7$. In this basis, the anti-canonical bundle of S is given by:

$$-k = 3e_0 - (e_1 + \dots + e_7).$$

The morphism $\phi_{(-k)} : S \rightarrow \mathbb{P}^2$ associated to the anti-canonical linear system is a double cover of the plane branched along a smooth plane quartic curve $Q(S)$. There are exactly 56 effective divisors in $Pic(S)$ with self-intersection -1 , given by the preimages of the 28 bitangents of $Q(S)$. Consider now the lattice:

$$R = k^\perp = \{x \in Pic(S) : (x, k) = 0\},$$

where $(,)$ denotes the intersection form on $Pic(S)$. This a root lattice of type E_7 and a root basis is given by:

$$\alpha_0 = e_0 - (e_1 + e_2 + e_3),$$

$$\alpha_i = e_i - e_{i+1}, \quad i = 1, \dots, 6.$$

We denote with $W(E_7)$ the associated Weyl group.

Let i be the covering involution of $\phi_{(-k)}$, then the induced involution i^* on $Pic(S)$ generates the center of the Weyl group $W(E_7)$ (see 4., Ch.VII, [4]). It is easy to see that

$$i^*(e_i) = -k - e_i, \quad i = 1, \dots, 7,$$

$$i^*(e_0) = -3k - e_0.$$

The eigenlattices $H_\pm$ of i^* relative to ± 1 are given by

$$H_+ = \langle k \rangle, \quad H_- = R.$$

Let X be a *K3 surface* i.e. a surface with $q(X) = 0$ and trivial canonical bundle. We denote with ω_X a generator for the vector space $H^{2,0}(X)$ of holomorphic two-forms on X , with $Pic(X)$ the Picard lattice and with $T(X)$ the transcendental lattice of X . We recall that the cohomology group $H^2(X, \mathbb{Z})$ is an even unimodular lattice isometric to $L_{K3} = U^{\oplus 3} \oplus E_8^{\oplus 2}$. The intersection form on L_{K3} is denoted with $(,)$.

2. THE MODEL BY KONDO

2.1. Geometric construction. We briefly recall the geometric construction introduced by Kondō in [7]. Let C be a smooth quartic curve in $\mathbb{P}^2$ defined by a homogeneous polynomial $f \in \mathbb{C}[x, y, z]$ of degree 4:

$$C = \{(x, y, z) \in \mathbb{P}^2 : f(x, y, z) = 0\}.$$

Consider the 4:1 cyclic cover of $\mathbb{P}^2$ branched along the divisor C :

$$\pi : X_C \xrightarrow{4:1} \mathbb{P}^2.$$

The surface X_C is a $K3$ surface with a degree four polarization, in coordinates it can be given by:

$$X_C = \{(x, y, z, t) \in \mathbb{P}^3 : t^4 = f(x, y, z)\}.$$

Let σ be a generator for the covering transformation group of π . We can assume that:

$$\sigma(x, y, z, t) = (x, y, z, it).$$

Let $\tau = \sigma^2$, then the morphism π factors naturally through the double cover of $\mathbb{P}^2$ branched along C :

$$\pi_1 : S_C \xrightarrow{2:1} \mathbb{P}^2,$$

where $S_C = X_C / \langle \tau \rangle$ is a Del Pezzo surface of degree 2 and $\pi_1 = \phi_{(-k)}$ is the morphism associated to the anti-canonical linear system of S_C . The geometry of the above construction is then described by the following commutative diagram:

$$\begin{array}{ccc} X_C & \xrightarrow{\pi_2} & S_C \\ \searrow \pi & & \swarrow \pi_1 \\ & \mathbb{P}^2 & \end{array}$$

The $K3$ surface X_C is L_+ -polarized, where L_+ denotes the pull-back of $\text{Pic}(S_C)$ and the isometry σ^* induced by σ on the cohomology lattice acts as a 4-th root of unity on the vector space of holomorphic two-forms. In other words, the period point of X_C belongs to $\mathbb{P}(W_{\mathbb{C}})$, where $W_{\mathbb{C}}$ is an eigenspace of σ^* in $L_+^{\perp} \otimes \mathbb{C}$. In fact, if $L_- = L_+^{\perp}$, then $W_{\mathbb{C}} \cong L_- \otimes_{\mathbb{Z}} \mathbb{R}$ and the action of σ^* gives to $W = L_- \otimes_{\mathbb{Z}} \mathbb{Q}$ the structure of a vector space over $k = \mathbb{Q}[i]$.

The vector space $W_{\mathbb{C}}$ is equipped with the hermitian form

$$\varphi(z, w) = (z, \bar{w})$$

of signature $(1, 6)$. The period domain $\mathcal{M}$ for polarized $K3$ surfaces (X_C, σ^*) is then an arithmetic quotient of the six dimensional complex ball B in $\mathbb{P}(W_{\mathbb{C}})$ defined by $\varphi(z, z) < 0$:

$$\mathcal{M} = B/\Gamma,$$

$$\Gamma = \{\gamma \in O(L_-) : \gamma \circ \sigma^* = \sigma^* \circ \gamma\} \cong U(\varphi) \cap M(7, \mathbb{Z}[i]).$$

The above construction defines a period map:

$$\mathcal{P} : \mathcal{M}_3 \setminus \mathcal{M}_3^h \longrightarrow \mathcal{M},$$

where $\mathcal{M}_3^h$ denotes the hyperelliptic locus. In fact, the image of the period map lies in the complement of two divisors $\mathcal{D}_n, \mathcal{D}_h$, called *mirrors*, corresponding to non-ample polarized $K3$ surfaces. The main result in [7] is:

Theorem 2.1 (S. Kondō, Theorem 2.5, [7]). *The period map gives an isomorphism:*

$$\mathcal{P} : \mathcal{M}_3 \setminus \mathcal{M}_3^h \longrightarrow \mathcal{M} \setminus (\mathcal{D}_n \cup \mathcal{D}_h).$$

2.2. Isometries. We now describe the actions of covering transformations on the cohomology lattice. Let τ^* be the involution on $H^2(S_C, \mathbb{Z})$ induced by τ , its eigenlattices are given by:

$$L_{\pm} = \{x \in H^2(X_C, \mathbb{Z}) : \tau^*(x) = \pm x\}.$$

In fact, the invariant lattice L_+ is the Picard lattice of the generic $K3$ surface X_C and $L_- = L_+^{\perp}$. The isomorphism classes of the eigenlattices can be easily computed:

Lemma 2.2.

$$L_+ \simeq \langle 2 \rangle \oplus A_1^{\oplus 7}, \quad L_- \simeq \langle 2 \rangle^{\oplus 2} \oplus D_4^{\oplus 3}.$$

Proof. From Theorem 4.2.2, [10] and Theorem 3.6.2, [11] it follows that L_+ is a 2-elementary lattice of signature $(1, 7)$ with $r(L_+) = \ell(L_+) = 8$ and $\delta(L_+) = 1$. Hence obviously: $r(L_-) = 14$, $\ell(L_-) = 8$ and $\delta(L_-) = 1$. By Theorem A.2 the isomorphism classes of the two lattices are determined uniquely by the set of invariants (s_+, s_-, ℓ, δ) . Hence it is enough to check that the lattices in the right hand sides have the same set of invariants, this follows from the remarks in the Appendix. For the proof of the first isomorphism see also §2, [7]. $\square$

From Lemma 2.2 it follows that $L_+ = \pi_2^* \text{Pic}(S_C)$. For $x \in \text{Pic}(S_C)$ we define:

$$\tilde{x} = \pi_2^*(x) \in L_+.$$

In particular, fixed a blowing-up morphism $b : S_C \longrightarrow \mathbb{P}^2$ with the notation in section 1, a basis for L_+ is given by:

$$\tilde{e}_0, \tilde{e}_1, \dots, \tilde{e}_7.$$

The action of σ^* on L_+ is clearly the pull-back along π_2 of the action of the involution i^* on $\text{Pic}(S)$ defined in the previous section. In particular, let $\tilde{R} = \pi_2^*(R)$ and $\tilde{H}_{\pm}$ be the eigenlattices of σ^* on L_+ . From the remarks in section 1 immediately follows:

Lemma 2.3.

$$\tilde{H}_+ = \langle \tilde{k} \rangle \cong \langle 4 \rangle, \quad \tilde{H}_- = \tilde{R} \cong E_7(2).$$

Moreover, the action of the Weyl group $W(E_7)$ lifts to L_+ giving a subgroup:

$$W(E_7) \subset O(L_+)$$

of isometries commuting with σ^* .

A natural basis for L_- is given by:

$$t_1, t_2, f_1^1, \dots, f_4^1, f_1^2, \dots, f_4^2, f_1^3, \dots, f_4^3,$$

where t_1, t_2 is a basis for $\langle 2 \rangle^{\oplus 2}$ and $f_1^i, \dots, f_4^i$ is a basis for the i -th copy of D_4 as in the Appendix. Then we have

Lemma 2.4. *There exists an isomorphism $L_- \cong \langle 2 \rangle^{\oplus 2} \oplus D_4^{\oplus 3}$ such that the action of the isometry σ^* on L_- preserves $\langle 2 \rangle^{\oplus 2}$ and each copy of D_4 . In fact it is given by the matrix $J_1 \oplus J_2^{\oplus 3}$ (see the Appendix) with respect to the previously defined basis of L_- .*

Proof. See [1]. □

The discriminant groups of the eigenlattices of τ^* are given by (Proposition A.1):

$$A_{L_+} \cong A_{L_-} \cong \mathbb{Z}_2^{\oplus 8}.$$

In fact, we consider the natural bases:

$$\begin{aligned} A_{L_+} &= \langle \tilde{e}_0/2, \tilde{e}_1/2, \dots, \tilde{e}_7/2 \rangle, \\ A_{L_-} &= \langle t_1/2, t_2/2, \alpha_1^i, \alpha_2^i, i = 1, 2, 3 \rangle, \end{aligned}$$

where α_1^i, α_2^i generate the discriminant group of the i -th copy of D_4 in L_- (their expression in terms of $f_1^i, \dots, f_4^i$ is given in the Appendix). By Proposition A.4, the primitive embedding of L_+ in L_{K3} is determined by an isomorphism

$$\gamma : A_{L_+} \longrightarrow A_{L_-} \quad \text{with} \quad q_{L_-} \circ \gamma = q_{L_+}.$$

Lemma 2.5. *The isomorphism γ is given by the matrix*

$$M(\gamma) = \begin{pmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

up to the action of $O(A_{L_+})$ (with respect to the previously defined bases).

Proof. The primitive embedding of L_+ in L_{K3} is unique by Theorem 1.14.4, [10]. Hence, by Proposition A.4, the isomorphism γ is unique up to the action of the image of the natural homomorphism

$$\psi : O(L_+) \longrightarrow O(A_{L_+}).$$

In fact, by Theorem 1.14.2 in [10], the homomorphism ψ is surjective. Hence, it is enough to check that $M(\gamma)$ defines an isomorphism preserving the quadratic forms. □

The following result describes all vectors in A_{L_+} with assigned self-intersection with respect to the quadratic form q_{L_+} and the action of σ^* on them:

Lemma 2.6 (Lemma 2.1, [7]). *The vectors in A_{L_+} can be divided in four classes accordingly to their self-intersection:*

$$\begin{aligned} C_1 &= \{x \in A_{L_+} \mid q_{L_+}(x) = -1/2\}, \\ C_2 &= \{x \in A_{L_+} \mid q_{L_+}(x) = 1/2\}, \\ C_3 &= \{x \in A_{L_+} \mid q_{L_+}(x) = 1\}, \\ C_4 &= \{x \in A_{L_+} \mid q_{L_+}(x) = 0\}. \end{aligned}$$

- An element in C_1 is represented uniquely by one of the following 56 vectors:

$$\begin{aligned} &\frac{\tilde{e}_i}{2}, \quad \frac{\tilde{k} - \tilde{e}_i}{2}, \quad 1 \leq i \leq 7, \\ &\frac{\tilde{e}_0 - \tilde{e}_i - \tilde{e}_j}{2}, \quad \frac{\tilde{k} - \tilde{e}_0 + \tilde{e}_i + \tilde{e}_j}{2}, \quad 1 \leq i < j \leq 7. \end{aligned}$$

- An element in C_2 is represented uniquely by one of the following 72 vectors:

$$\frac{\tilde{e}_0}{2}, \frac{\tilde{k} - \tilde{e}_0}{2},$$

$$\frac{\tilde{e}_i + \tilde{e}_j + \tilde{e}_k}{2}, \frac{\tilde{k} - \tilde{e}_i - \tilde{e}_j - \tilde{e}_k}{2}, \quad 1 \leq i < j < k \leq 7.$$

- An element in C_3 is represented uniquely by one of the following 64 vectors:

$$\frac{\tilde{k}}{2}, \frac{\tilde{e}_i - \tilde{e}_j}{2}, \quad 1 \leq i < j \leq 7,$$

$$\frac{\tilde{e}_0 - \tilde{e}_i - \tilde{e}_j - \tilde{e}_k}{2}, \quad 1 \leq i < j < k \leq 7,$$

$$\frac{2\tilde{e}_0 - \sum_i \tilde{e}_i + \tilde{e}_j}{2}, \quad 1 \leq j \leq 7.$$

- An element in C_4 is represented uniquely by one of the following 64 vectors:

$$0, \frac{\tilde{k} + \tilde{e}_i - \tilde{e}_j}{2}, \quad 1 \leq i < j \leq 7,$$

$$\frac{\tilde{k} + \tilde{e}_0 - \tilde{e}_i - \tilde{e}_j - \tilde{e}_k}{2}, \quad 1 \leq i < j < k \leq 7,$$

$$\frac{\tilde{k} + 2\tilde{e}_0 - \sum_i \tilde{e}_i + \tilde{e}_j}{2}, \quad 1 \leq j \leq 7.$$

The isometry σ^* acts as:

$$\sigma^*(x) = \begin{cases} \tilde{k}/2 - x & \text{if } x \in C_1 \cup C_2 \\ x & \text{if } x \in C_3 \cup C_4. \end{cases}$$

Let

$$N_+ = \{x \in L_+^* : q_{L_+}(x) \in \mathbb{Z}/2\mathbb{Z}\}.$$

Notice that:

$$N_+/L_+ = \langle \tilde{k}/2, \tilde{\alpha}_1/2, \dots, \tilde{\alpha}_6/2 \rangle.$$

We denote with $O(A_{L_+})$ the group of automorphisms of A_{L_+} respecting the discriminant quadratic form q_{L_+} . Then we have

Lemma 2.7. *The natural homomorphism $W(E_7) \longrightarrow O(A_{L_+})$ is surjective. Moreover, the action of $O(A_{L_+})$ on vectors in Lemma 2.6 is transitive on classes C_1 and C_2 and has two orbits on classes C_3 and C_4 .*

Proof. The first assertion follows from Lemma 2.3, [7]. The elements of C_1 are of the form $\tilde{x}/2$ where x is the class of a (-1) -curve on the Del Pezzo surface S . The Weyl group $W(E_7)$ acts transitively on (-1) -curves (see for example Lemma 4, Ch. V, [4]), hence the action of $O(A_{L_+})$ is transitive on C_1 .

Consider now the elements in the class C_2 :

$$\tilde{e}_0/2, v_{ijk} = (\tilde{e}_i + \tilde{e}_j + \tilde{e}_k)/2.$$

Let $\alpha_{ijk} = \tilde{e}_0 - v_{ijk}$, then:

$$s_{\alpha_{ijk}}(v_{ijk}) = 3\tilde{e}_0 - 2v_{ijk}/2 = \tilde{e}_0/2 \pmod{L_+}.$$

Similarly, two vectors v_{ijk} and $v_{i'j'k'}$ are interchanged by the isometry obtained as composition of the three reflections with respect to $e_i - e_{i'}$, $e_j - e_{j'}$, $e_k - e_{k'}$. This gives the assertion for C_2 .

The element $\tilde{k}/2$ in the class C_3 is preserved by the Weyl group. The other 63 classes are of the form $\tilde{x}/2$ where x is a (positive) root of E_7 . Since $W(E_7)$ acts transitively on roots (see for example [4]), the action is transitive on C_3 . The elements in C_4 are obtained from those in C_3 by adding $\tilde{k}/2$, hence the assertion follows easily. $\square$

3. HEEGNER DIVISORS AND EMBEDDINGS

Definition 3.1. Let $r \in L_-$ be a primitive vector and let H_r be the hyperplane section of B orthogonal to r :

$$H_r = \{z \in B : (z, r) = 0\}.$$

The *Heegner divisor* $\mathcal{D}_r$ associated to r is the image of H_r in $\mathcal{M}$. A Heegner divisor $\mathcal{D}_r$ is of *type* n if $r^2 = -2n$.

Remark 3.2.

i) For $r \in L_-$ let $z(r) = r - i\sigma^*(r)$, then $z(r) \in W$ and

$$H_r = \{z \in W : \varphi(z, z(r)) = 0\},$$

where φ is the hermitian form on W defined in section 2. This implies that a Heegner divisor can be equivalently defined as the image in $\mathcal{M}$ of a hyperplane section of B defined over $k = \mathbb{Q}[i]$. In the language of [8], Heegner divisors are quotients of an arithmetically defined arrangement of hyperplanes in the ball.

ii) It can be easily seen that $H_r = H_{\sigma^*(r)}$ and $H_r = \emptyset$ if $n \leq 0$.

With the notation in the Appendix:

$$\Lambda_r = \langle r, \sigma^*(r) \rangle \cong A_1(n)^{\oplus 2}.$$

Hence, giving a Heegner divisor of type $n \geq 1$ in $\mathcal{M}$ is equivalent to assign a primitive embedding of the lattice $A_1(n)^{\oplus 2}$ in L_- up to the action of the group Γ . Given such an embedding, let $\Lambda_r^\perp$ be the orthogonal complement of Λ_r in L_- and P be the orthogonal complement of $\Lambda_r^\perp$ in L_{K3} . We have an embedding

$$L_+ \oplus \Lambda_r \subset P$$

with finite quotient group

$$M_r = P/(L_+ \oplus \Lambda_r).$$

We start describing the structure of the group M_r .

Proposition 3.3. *Let $r \in L_-$ and M_r as above then:*

- a) *if n is even then $M_r \cong \mathbb{Z}_2$ and $(r, L_-) = \mathbb{Z}$;*
- b) *if n is odd then:*

$$M_r \cong \begin{cases} \mathbb{Z}_2 \oplus \mathbb{Z}_2 & \text{if } (r, L_-) = 2\mathbb{Z}, \\ \mathbb{Z}_2 & \text{if } (r, L_-) = \mathbb{Z}. \end{cases}$$

Proof. The group M_r has a natural embedding in $A_{L_+} \oplus A_{\Lambda_r}$. Moreover, since the embedding of Λ_r in P is primitive, the projections on the two factors are isomorphisms (see Appendix):

$$M_r \cong M_1 \subset A_{L_+}, \quad M_r \cong M_2 \subset A_{\Lambda_r}$$

such that

$$q_{L_+|M_1} = -q_{\Lambda_r|M_2}.$$

Since

$$A_{L_+} \cong \mathbb{Z}_2^{\oplus 8}, \quad A_{\Lambda_r} \cong \mathbb{Z}_{2n}^{\oplus 2}$$

it follows that:

$$M_r \cong \mathbb{Z}_2^{\oplus m_r}, \quad 0 \leq m_r \leq 2.$$

We first prove that the case $m_r = 0$ does not appear i.e. $L_+ \oplus \Lambda_r$ can not be primitively embedded in L_{K3} . This means that for any embedding $L_+ \oplus \Lambda_r \hookrightarrow L_{K3}$ there is a primitive element $(x, y) \in L_+ \oplus \Lambda_r$ such that $(x, y)/2 \in L_{K3}$. For this, it suffices to prove that, for some $y \in \Lambda_r$, $y/2$ is not trivial in A_{L_-} , since in this case (see § 2.2):

$$\gamma^{-1}(y/2) + y/2 \in L_{K3}.$$

Let $r = (t, d_1, d_2, d_3) \in L_-$ be a primitive vector with $t \in \langle 2 \rangle^{\oplus 2}$ and $d_i \in D_4$, $i = 1, 2, 3$ according to the direct sum decomposition of L_- given in Lemma 2.2. From the remarks in the Appendix (in particular Remark A.3) it follows that we have these cases:

- i) d_1, d_2, d_3 are divisible by two, then t is not divisible by two, hence $r/2 = t/2$ is not trivial in A_{L_-} ;
 - ii) at least one among d_1, d_2, d_3 is not divisible by two and not all its coefficients are odd, then $r + \sigma^*(r)/2$ is non-trivial in A_{L_-} ;
 - iii) at least one among d_1, d_2, d_3 is not divisible by two and the non-two divisible d'_i s all have odd coefficients, then $r/2$ is in L_-^* and it is non-trivial in A_{L_-} .
- Hence $m_r \in \{1, 2\}$.

We now characterize the remaining cases. According to the previous discussion, there are two possibilities:

- i) $r/2 \in A_{L_-}$ i.e. $(r, L_-) = 2\mathbb{Z}$, then

$$M_r = \langle r/2, \sigma^*(r)/2 \rangle.$$

Since Λ_r is primitive in L_- , we have that $\sigma^*(r)/2 \neq r/2$ in A_{L_-} . In particular n is not even, since otherwise $q(r/2) = -n/2 \in \mathbb{Z}$ hence $\sigma^*(r)/2 = r/2$ in A_{L_-} by Lemma 2.6). Therefore $m_r = 2$.

- ii) $r/2 \notin A_{L_-}$ i.e. $(r, L_-) = \mathbb{Z}$, then $m_r = 1$ and

$$M_r = \langle (r + \sigma^*(r))/2 \rangle.$$

□

We call m_r the *index* of the Heegner divisor $\mathcal{D}_r$, $r \in L_-$.

Proposition 3.4 (Existence).

- i) *There exist Heegner divisors of type n and index 1 for every $n \geq 1$;*
 - ii) *there exist Heegner divisors of type n and index 2 for every odd $n \geq 1$.*
- (Note that the case of even type and index 2 is excluded by Proposition 3.3).

Proof. Consider the following vectors in L_-

$$\begin{aligned} r_1(1) &= f_1^1 + f_1^2, \quad r_1(k) = (k-1)t_1 + kf_1^1 + f_1^2 \quad (k > 1), \\ r_2(k) &= kt_1 + (k+1)f_1^1, \\ r_3(k) &= (k-1)t_1 + kf_1^1 + f_1^2 + f_1^3, \\ r_4(k) &= t_1 + 2kt_2 + (k+1)(f_1^1 + f_1^2) + k(f_1^2 + f_2^2), \end{aligned}$$

with respect to the basis of L_- defined in section 2. It is easy to check that the rank two lattice $\Lambda_i(k) = \langle r_i(k), \sigma^*(r_i(k)) \rangle$ is primitive in L_- , since the rank two

minors of the matrix with rows $r_i(k), \sigma^*(r_i(k))$ have no common factor. Moreover, notice that

$$r_i(k)^2 = \begin{cases} -2(2k) & \text{if } i = 1, \\ -2(2k+1) & \text{if } i = 2, 3, \\ -2(4k+1) & \text{if } i = 4, \end{cases}$$

in particular the Heegner divisor $\mathcal{D}_{r_i(k)}$ has even type $n = 2k$ for $i = 1$ and odd type otherwise. By Proposition 3.3 the Heegner divisor $\mathcal{D}_{r_i(k)}$ has index 1 if $i = 1, 3$ since $(r_i(k), L_-) = \mathbb{Z}$, hence i) is proved.

For $i = 2$ it can be easily checked that the index is 2 iff k is odd i.e. the type $n = 2k + 1 \equiv 3 \pmod{4}$. For $i = 4$ the type is $n = 4k + 1 \equiv 1 \pmod{4}$ and the index is 2 by Proposition 3.3, hence ii) is proved. $\square$

The following remark will be useful in the next section

Lemma 3.5. *Let $\mathcal{D}_r$ be a Heegner divisor of even type, then $\tilde{k}/2 \notin M_r$.*

Proof. Since $\mathcal{D}_r$ has index 1 by Proposition 3.3 we have $M_r = \langle (r + \sigma^*(r))/2 \rangle$. From Lemma 2.5 it follows that the isomorphism $\gamma : A_{L_+} \rightarrow A_{L_-}$ sends $\tilde{k}/2$ to

$$\gamma(\tilde{k}/2) = (1/2, 1/2, 0, \dots, 0)$$

(in the usual coordinates for L_-). Consider a vector:

$$r = (h_1, h_2, a_1, \dots, a_4, b_1, \dots, b_4, c_1, \dots, c_4),$$

such that Λ_r is primitive in L_- and assume that $(r + \sigma^*(r))/2 = \gamma(\tilde{k}/2)$ in A_{L_-} . This means that:

- i) h_1 and h_2 are odd,
- ii) a_3, b_3 and c_3 are even,
- iii) $a_1 + a_2 + a_4, b_1 + b_2 + b_4$ and $c_1 + c_2 + c_4$ are even.

In this case it can be easily checked that $(r, L_-) = 2\mathbb{Z}$, giving a contradiction by Proposition 3.3. $\square$

4. HEEGNER DIVISORS AND GENUS 3 CURVES

Let X_r be the generic $K3$ surface with period point in a Heegner divisor $\mathcal{D}_r$, then its Picard number is equal to 10 and

$$L_+ \oplus \Lambda_r \subset \text{Pic}(X_r).$$

In this section we describe the loci of genus three curves corresponding to Heegner divisors of given type and index. We first recall the special role of Heegner divisors of type 1 in Kondō's construction.

4.1. Type 1.

Lemma 4.1. *A period point in $\mathcal{M}$ belongs to a Heegner divisor of type 1 iff the corresponding $K3$ surface is not the four cyclic cover of the plane branched along a smooth plane quartic.*

Proof. See Theorem 2.5, [7]. $\square$

Proposition 4.2. *The mirrors $\mathcal{D}_n, \mathcal{D}_h$ are the unique Heegner divisors of type 1:*

- $\mathcal{D}_n$ has index 1 and its generic point corresponds to a plane quartic with a node;

- $\mathcal{D}_h$ has index 2 and its generic point corresponds to a smooth hyperelliptic genus three curve.

Proof. By Lemma 3.3, Theorem 4.3 and Theorem 5.4 in [7]. $\square$

Remark 4.3.

- i) In [1] it is proved that the period map $\mathcal{P}$ can be defined on the blow-up of the GIT moduli space of semistable plane quartics in one point. This period map gives an isomorphism between $\mathcal{D}_n$ and the locus of stable singular quartics. Besides, it is proved that the Baily-Borel compactification of $\mathcal{D}_h$ is isomorphic to the GIT moduli space of semistable sets of eight unordered points in $\mathbb{P}^1$ (see also [6]).
- ii) The transcendental lattices of the generic $K3$ surface in $\mathcal{D}_n$ and $\mathcal{D}_h$ are given respectively by:

$$\begin{aligned} T_n &\cong U^{\oplus 2} \oplus A_1^{\oplus 8}, \\ T_h &\cong U(2)^{\oplus 2} \oplus D_8. \end{aligned}$$

4.2. Type $n > 1$.

Definition 4.4. Let C be a plane quartic and $\pi : X_C \rightarrow \mathbb{P}^2$ be the four cyclic cover of the plane branched along C . An irreducible plane curve D is a *splitting curve* for C if the inverse image $\pi^{-1}(D)$ is the union of four distinct curves.

It is easy to see that a smooth plane curve D is a splitting curve for C if and only if the restricted cover $\pi|_{\pi^{-1}(D)}$ is trivial. If the curve is not smooth it can happen that it is partially normalized by the cover π i.e. the four curves in $\pi^{-1}(D)$ are not isomorphic to D and they can intersect each other outside the ramification curve. We will see examples of this behaviour in the last section.

With X_r as before, we call $\pi : X_r \rightarrow \mathbb{P}^2$ the four cyclic cover of $\mathbb{P}^2$, C_r the corresponding branch quartic and $\pi_2 : X_r \rightarrow S_r$ the associated double cover of the Del Pezzo surface S_r (see section 2). We start with a simple remark

Lemma 4.5. *Let $\mathcal{D}_r$ be a Heegner divisor of type $n > 1$, then C_r admits a splitting curve of even degree.*

Proof. As noticed before, we have $L_+ \oplus \Lambda_r \subset \text{Pic}(X_r)$. Consider a class of the form $w = \tilde{x} + r$ where $\tilde{x} \in L_+$ with $\tilde{x}^2 \geq 2n - 2$. Then $w^2 \geq -2$, hence w or $-w$ is effective by the Riemann Roch Theorem. Assume that w is effective and write $w = \sum a_i w_i$, where a_i are positive integers and w_i are irreducible curves. Then we can assume that $w_1 = \tilde{x}_1 + r_1$ with $r_1 \neq 0$. The σ^* -orbit of w_1 clearly contains four distinct elements: $w_1, \sigma^*(w_1), \tau^*(w_1) = \tilde{x}_1 - r_1, (\sigma^*)^3(w_1) = \sigma^*(\tilde{x}_1) - \sigma^*(w_1)$. Hence the image $\pi(R)$ of a curve R in the class w_1 is a splitting curve for C . The degree of R is given by $\deg(R) = (\tilde{x}_1, -\tilde{k}) = 2(x_1, -k)$. $\square$

Theorem 4.6. *Let $\mathcal{D}_r$ be a Heegner divisor of type $n > 1$ (so $r \in L_-$, $r^2 = -2n$) with index $m_r \in \{1, 2\}$. Then there exists a rational splitting curve D for C_r with the following properties:*

1)

$$\deg(D) = \begin{cases} 2(n-1) & \text{if } m_r = 1, \\ n-2 & \text{if } m_r = 2. \end{cases}$$

2) *The inverse image $\pi_1^{-1}(D)$ in S_r is the union of two smooth rational curves (hence $\pi^{-1}(D)$ is the union of four smooth rational curves in X_r).*

Proof. The idea is to construct the class of a (-2) -curve in $Pic(X_r) \setminus L_+ \oplus \Lambda_r$ starting from the information given by the structure of the group M_r .

i) If $m_r = 1$ (i.e. $M_r \cong \mathbb{Z}_2$) then there exists $\tilde{x} \in L_+$ such that:

$$(\tilde{x} + f)/2 \in Pic(X_r)$$

where $f = r + \sigma^*(r)$. Notice that $(\tilde{x}' + f)/2 \in Pic(X_r)$ for any $\tilde{x}' \in L_+^*$ with $\tilde{x}'/2 = \tilde{x}/2$ in A_{L_+} . Moreover

$$q(\tilde{x}/2) = -q(f/2) = n.$$

Then $\tilde{x}/2 \in A_{L_+}$ belongs to the class C_3 in Lemma 2.6 if n is odd and to the class C_4 if n is even.

If n is odd we can assume (up to the action of the Weyl group, see Lemma 2.7 and Lemma 3.5) that:

$$\tilde{x}/2 = (\tilde{e}_1 - \tilde{e}_2)/2 \in A_{L_+}.$$

Let $x' = (n-1)e_0 - (n-2)e_1 - e_2 \in Pic(S_r)$, then $\tilde{x}'/2 = \tilde{x}/2$ in A_{L_+} . Hence by a previous remark

$$y = (\tilde{x}' + f)/2 \in Pic(X_r)$$

and $y^2 = (x')^2/2 - n = -2$. Notice that $y + \tau^*(y) = x'$ and the arithmetic genus $p_a(x')$ is zero by the adjunction formula.

The class y can be written as $y = \sum_{i=1}^l a_i y_i$, where a_i are positive integers and y_i are irreducible curves. Since $p_a(x') = 0$, there exists an irreducible component y_1 of y which is a smooth rational curve and doesn't belong to L_+ . This can be written as

$$y_1 = (\tilde{x}_1 + f_1)/2, \quad \tilde{x}_1 \in L_+, \quad f_1 \in \Lambda_r, \quad f_1 \neq 0.$$

Since $y^2 = y_1^2 = -2$ and $f_1^2 \leq f^2$ we have $x_1^2 \geq (x')^2 = 2(n-2)$. Moreover, since x_1 is a component of x' , its arithmetic genus is zero. This gives

$$0 = p_a(x_1) = \frac{x_1^2 + (x_1, k)}{2} + 1 \geq \frac{2(n-2) + (x_1, k)}{2} + 1,$$

hence

$$(y_1, -k) = (x_1, -k) \geq 2(n-1) = (y, -k).$$

This implies that $y_1 = y$ i.e. y is a (-2) -curve. Let R be the curve in the class y , then the plane curve $D = \pi(R)$ is a rational splitting curve for C_r and:

$$\deg(D) = -(x', k) = 2(n-1).$$

If n is even, then we can assume (up to the action of the Weyl group, see Lemma 2.7) that:

$$\tilde{x}/2 = (\tilde{k} + 2\tilde{e}_0 - \sum_i \tilde{e}_i + \tilde{e}_1)/2 = (\tilde{e}_0 - \tilde{e}_1)/2 \in A_{L_+}.$$

The proof follows as in the odd case with the choice $x' = (n-1)e_0 - e_1 - (n-2)e_2$.

ii) If $m_r = 2$ (i.e. $M_r \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$) then there exists $\tilde{x} \in L_+$ such that:

$$(\tilde{x} + r)/2, \quad \sigma^*(\tilde{x} + r)/2 \in Pic(X_r),$$

where

$$q(\tilde{x}/2) = -q(r/2) = n/2.$$

Then $\tilde{x}/2 \in A_{L_+}$ belongs to the class C_1 in Lemma 2.6 if $n \equiv 3 \pmod{4}$ and to the class C_2 if $n \equiv 1 \pmod{4}$.

If $n \equiv 3 \pmod{4}$, then we can assume that:

$$\tilde{x}/2 = \tilde{e}_1/2.$$

Let $n = 4s + 3$ and consider the vector $x' = 2se_0 - (2s - 1)e_1$. Note that $p_a(x') = 0$ by the adjunction formula. Then

$$y = (x' + r)/2 \in \text{Pic}(X_r)$$

and $y^2 = -2$. It can be proved as in i) that y is a smooth (-2) -curve and if R is a curve representing y , then $D = \pi(R)$ is a rational splitting curve with

$$\deg(D) = -(x', -k) = n - 2.$$

If $n \equiv 1 \pmod{4}$, then we can assume that:

$$\tilde{x}/2 = \tilde{e}_0/2.$$

Let $n = 4s + 1$ and consider the class $x' = (s - 1)(e_0 - e_1)$. Defining y as above, similar computations give a rational splitting curve D with:

$$\deg(D) = -(x', k) = n - 2.$$

□

Remark 4.7. Note that, in the proof of Theorem 4.6, (the classes of) the preimages of the splitting curves in the Del Pezzo surface are given explicitly up to the action of the Weyl group.

Corollary 4.8. *The Picard lattice of the K3 surface X_r is generated by L_+ and two smooth rational curves $R, \sigma^*(R)$ with $R \notin L_+ \oplus \Lambda_r$.*

Proof. Obviously $\text{Pic}(X_r) \supset \langle L_+, R, \sigma^*(R) \rangle$. Let $R = (\tilde{x} + y)/2$, where $\tilde{x} \in L_+$ and $y \in \Lambda_r$ are primitive. Then the thesis follows since the embedding of L_+ in $\text{Pic}(X_r)$ is primitive and the classes $R, \sigma^*(R)$ are not divisible in L_{K3} . □

We also give a partial converse to the previous result:

Proposition 4.9. *Let C be a plane quartic admitting a rational splitting curve D such that the inverse image of D in S_C is the union of two smooth rational curves with primitive classes in $\text{Pic}(S_C)$. Let d be the minimal degree of a curve D with this property. Then X_C belongs to a Heegner divisor*

- i) of index $m = 2$ and type $n = d + 2$ if d is odd;*
- ii) of index $m = 1$ and type $n = (d + 2)/2$ if d is even.*

Proof. Let R be one of the inverse images of D in X_C . Notice that:

$$R = (x + y)/2 \in \text{Pic}(X_C),$$

where

$$x = R + \tau^*(R) \in L_+, \quad y = R - \tau^*(R) \in L_- \cap \text{Pic}(X_C).$$

An easy computation gives:

$$x^2 = 2(d - 2), \quad y^2 = -2(d + 2).$$

Notice that $y \neq 0$ in L_- since otherwise $x = 2R$ (which is the class of a preimage of D in S_C) would be divisible in L_+ . Thus X_C belongs to a Heegner divisor $\mathcal{D}_r$ with $y \in \Lambda_r$. Let $\Lambda_r^\perp$ be the orthogonal complement of Λ_r in L_- and P_r be the orthogonal complement of $\Lambda_r^\perp$ in L_{K3} . Then we define

$$M_r = P_r/L_+ \oplus \Lambda_r.$$

Notice that:

$$\sum_{i=0}^3 (\sigma^*)^i(R) = -d\tilde{k}.$$

Then we have the following cases:

i) If d is odd, then:

$$\sum_{i=0}^3 (\sigma^*)^i(R)/2 = (x + \sigma^*(x))/2 = \tilde{k}/2 \neq 0 \in A_{L_+}.$$

Hence $x/2 \neq \sigma^*(x)/2 \in A_{L_+}$. Therefore

$$M_r \cong \langle x/2, \sigma^*(x)/2 \rangle \subset A_{L_+},$$

this means that $\mathcal{D}_r$ has index $m_r = 2$. By Theorem 4.6 (see also the proof) the curve C admits a rational splitting curve D' of degree $d' = -r^2/2 - 2$ with smooth and primitive preimages in the Del Pezzo surface such that $\pi^*(D') = \sum_{i=1}^4 (\sigma^*)^i(R')$, $i = 1, \dots, 4$ with $R' - \tau^*(R') = r$. Then:

$$(R' - \tau^*(R'))^2 = -2(d' + 2)$$

is maximal i.e. d' is minimal. Then $d = d'$ and $\mathcal{D}_r$ has type $n = d + 2$.

ii) If d is even, then:

$$\sum_i (\sigma^*)^i(R)/2 = (x + \sigma^*(x))/2 = 0 \in A_{L_+}.$$

Hence:

$$x/2 = \sigma^*(x)/2 \in A_{L_+}.$$

The case $x/2 = 0$ in A_{L_+} (i.e. $x = 2x' \in L_+$) can be excluded since otherwise the inverse images of D in S_C would belong to divisible classes. Then the Heegner divisor $\mathcal{D}_r$ has index $m_r = 1$:

$$M_r = \langle x/2 \rangle \subset A_{L_+}.$$

By applying Theorem 4.6 as in i), the minimality condition on the degree gives that $\mathcal{D}_r$ has type $n = (d + 2)/2$. $\square$

Corollary 4.10. *A plane quartic C admits a splitting curve of odd degree if and only if X_C belongs to a Heegner divisor of index 2.*

Proof. The result follows from Theorem 4.6 and the proof of Proposition 4.9 if one notices that in the odd degree case the inverse images of the splitting curve in $\text{Pic}(S_C)$ always have primitive classes. $\square$

Remark 4.11.

i) Splitting curves of even degree do not always give rise to Heegner divisors. From the proof of Proposition 4.9 it follows that this happens iff a inverse image R of the splitting curve in the $K3$ surface satisfies:

$$R - \tau^*(R) = 0.$$

For example, let C be the generic plane quartic with equation $f_4(x, y, z) = 0$. It is easy to prove that a plane quartic D of the form $l(x, y, z)^4 = f_4(x, y, z)$, where l is linear, is a splitting curve for C . The inverse image of D in X_C splits in four genus three curves $(\sigma^*)^i(R)$, $i = 0, \dots, 3$ which are hyperplane sections of X_C in $\mathbb{P}^3$ (i.e. they all belong to the class $-\tilde{k} \in L_+$).

- ii) If the splitting curve has even degree $d \not\equiv 2 \pmod{4}$ an easy computation shows that its inverse images have primitive classes in the Del Pezzo surface.
- iii) Theorem 4.6 and Proposition 4.9 describe the loci of plane quartics corresponding to the union of all Heegner divisors of given type and index. However, it is not clear if these loci are irreducible. In other words, it is unknown if an Heegner divisor is determined uniquely by its type and index. Note that this problem is equivalent to classify the embeddings of the lattice Λ_n in L_- up to the action of Γ .
- iv) In [6] Heegner divisors in $\mathcal{D}_h$ (in fact, this is a union of our Heegner divisors with fixed type and index intersected with $\mathcal{D}_h$) are introduced and the existence of an automorphic form vanishing exactly on them is proved.

5. EXAMPLES

In this section we give some examples of Heegner divisors corresponding to plane quartics with splitting curves of degree one, two and three. We will usually identify Heegner divisors in $\mathcal{M}$ and their inverse image by $\mathcal{P}$ in $\mathcal{M}_3$.

Let $C \subset \mathbb{P}^2$ be a plane curve of degree dd' with $d = 2, 4$ and let $\pi_d : Z \rightarrow \mathbb{P}^2$ be the $d:1$ cover of $\mathbb{P}^2$ branched along C . We start recalling the following result:

Proposition 5.1 (Proposition 1.7, Ch.3, [12]). *Let $D \subset \mathbb{P}^2$ be a curve not containing components of C . Let $\tilde{D} = \pi_d^{-1}(D)$. The restriction of the cover:*

$$\pi_{d|\tilde{D}} : \tilde{D} \longrightarrow D$$

is trivial if and only if there exists a curve $B \subset \mathbb{P}^2$ of degree d' such that:

$$C \cdot D = dB \cdot D.$$

By the remark after Definition 5.1, this result characterizes smooth splitting curves of C .

5.1. Quartics with a hyperflex. Let C be a generic smooth plane quartic with a hyperflex line L i.e.

$$L \cdot C = 4p, \quad p \in C.$$

After a projective transformation we can assume that the line L is given by the equation $x = 0$ and that the quartic C is of the form:

$$(1) \quad C : xf_3(x, y, z) + z^4 = 0,$$

where f_3 is a cubic polynomial in $\mathbb{C}[x, y, z]$. It is known that the locus of plane quartics with at least one hyperflex line is an irreducible closed subvariety of codimension one in the moduli space of genus three curves (see Proposition 4.9, Ch.I, [12]). As usual, let $\pi_1 : S_C \rightarrow \mathbb{P}^2$ and $\pi : X_C \rightarrow \mathbb{P}^2$ be the double and the 4:1 cover of $\mathbb{P}^2$ branched along C respectively.

Proposition 5.2. *The locus of plane quartics with a hyperflex line is the unique Heegner divisor of type 3 and index 2.*

Proof. Let C be a plane quartic with a hyperflex line L . By Proposition 5.1 both π_1 and π are trivial over L . Hence the line L is a splitting curve for C , moreover it splits in the union of two smooth rational curves on the Del Pezzo surface S_C . By Proposition 4.9 it follows that X_C belongs to a Heegner divisor of type $n = 3$ and index 2 (the degree of L is obviously minimal). The converse follows from Theorem 4.6. Since the locus of plane quartics with a hyperflex line is irreducible in $\mathcal{M}_3$,

there is only one Heegner divisor of type 3 and index 2. $\square$

We denote this Heegner divisor with $\mathcal{D}_{flex}$. We now compute the Picard lattice of the generic K3 surface X in $\mathcal{D}_{flex}$. Let M, M' be the inverse images of L in the Del Pezzo surface S .

$$\pi_1^*(L) = M + M'.$$

Each of them splits in the union of two smooth rational curves on X :

$$\pi_2^*(M) = M_1 + M_2, \quad \pi_2^*(M') = M'_1 + M'_2.$$

Notice that $\sigma^*(M_i) = M'_i$, $i = 1, 2$, in particular:

$$\begin{aligned} M_1 + M_2, \quad M'_1 + M'_2 &\in L_+, \\ M_1 - M_2, \quad M'_1 - M'_2 &\in L_- \cap \text{Pic}(X). \end{aligned}$$

Let $r = M_1 - M_2$, it is easy to check that $r^2 = -6$. Hence the period point of the K3 surface X lies in the hyperplane section of B determined by the vector $r \in L_-$. By Corollary 4.8 we have:

$$\text{Pic}(X) = \langle L_+, M_1, M'_1 \rangle.$$

Notice that M, M' are two (-1) -curves on S with:

$$M + M' = \pi_1^*(L) = -k.$$

Hence we can assume, up to the action of the Weyl group, that:

$$M = e_7, \quad M' = 3e_0 - (e_1 + \cdots + e_6 + 2e_7).$$

Then:

$$M_1 + M_2 = \tilde{e}_7, \quad M'_1 + M'_2 = 3\tilde{e}_0 - (\tilde{e}_1 + \cdots + \tilde{e}_6 + 2\tilde{e}_7).$$

The intersection matrix of the Picard lattice with respect to the basis $\tilde{e}_0, \tilde{e}_1, \dots, \tilde{e}_7, M_1, M'_1$ is then given by:

$$Q(X) = \begin{pmatrix} 2 & O_{1,6} & 0 & 0 & 3 \\ O_{6,1} & -2I_6 & O_6 & O_6 & I_{6,1} \\ 0 & 0 & -2 & -1 & 2 \\ 0 & 0 & -1 & -2 & 1 \\ 3 & 1 & 2 & 2 & 1 \end{pmatrix}$$

where $O_{1,6}$ and $O_{6,1}$ are respectively a row and a column zero vector of length 6, O_6 is the zero matrix of order 6, I_6 is the identity matrix of order 6 and $I_{6,1}$ a column vector of 1's.

5.2. Quartics with splitting conics. We consider the locus of smooth plane quartics C such that there exists an irreducible conic T with intersection divisor of the form:

$$T \cdot C = 4p + 4q,$$

where $p \neq q$. After a projective transformation we can assume that the conic T is given by:

$$T : xy - z^2 = 0$$

and that $p = (0, 1, 0)$, $q = (1, 0, 0)$. Then, up to projectivities, the equation of the quartic C is of the form:

$$C : (xy - z^2)f_2(x, y, z) + z^4 = 0,$$

where f_2 is a quadratic polynomial in $\mathbb{C}[x, y, z]$. It is known (see [5]) that a smooth plane quartic C has 63 one dimensional families of tangent conics and that each family contains six reducible conics (i.e. the union of two bitangents). We now prove that

Lemma 5.3. *The locus of plane quartics with an irreducible splitting conic is a codimension one closed irreducible subvariety of $\mathcal{M}_3$.*

Proof. Let $\Delta \subset \mathbb{P}^{14}$ be the discriminant locus corresponding to singular plane quartics. Notice that polynomials like the one defining C define a 6 dimensional irreducible subvariety A of $\mathbb{P}^{14} \setminus \Delta$. Let $\phi : \mathbb{P}^{14} \setminus \Delta \rightarrow \mathcal{M}_3 \setminus \mathcal{M}_3^h$ be the natural morphism to the moduli space of non-hyperelliptic genus three curves. This is a closed and surjective morphism (see Proposition 4.7, [12]). In particular, $V = \phi(A)$ is an irreducible closed subvariety of $\mathcal{M}_3 \setminus \mathcal{M}_3^h$. Note that the dimension of the fibers of ϕ over V is equal to one. In fact, it equals the dimension of:

$$\{g \in PGL(3, \mathbb{C}) : g(p) = p, g(q) = q, g(T) = T\}.$$

Thus V has codimension one in $\mathcal{M}_3 \setminus \mathcal{M}_3^h$. In fact, it can be proved as in [12] that V is also closed in $\mathcal{M}_3$. $\square$

Proposition 5.4. *The locus of plane quartics with an irreducible splitting conic is the unique Heegner divisor of type 2 and index 1.*

Proof. By Proposition 5.1, the inverse image of the conic T splits in the union of two smooth rational curves R, R' on the Del Pezzo surface and in four smooth rational curves $R_1, R_2, R'_1 = \sigma^*(R_1), R'_2 = \sigma^*(R_2)$ on the $K3$ surface X_C . By [5], R is equivalent to the union of two (-1) -curves L_1, M_1 with $(L_1, M_1) = 1$ and mapping to distinct bitangents of C . Notice that $(L_1, \sigma^*(M_1)) = 0$. We can assume, up to the action of the Weyl group, that:

$$L_1 = e_0 - e_1 - e_2, \quad \sigma^*(M_1) = 2e_0 - e_1 - e_2 - e_5 - e_6 - e_7.$$

Then

$$\begin{aligned} R &= L_1 + M_1 = 2e_0 - (e_1 + \cdots + e_4), \\ R' &= \sigma^*(L_1 + M_1) = 4e_0 - (e_1 + \cdots + e_4) - 2(e_5 + \cdots + e_7). \end{aligned}$$

In particular, R is a primitive class in $Pic(S_C)$. Then, by Proposition 4.9, X_C belongs to a Heegner divisor of type 2 and index 1. The converse follows from Theorem 4.6 and Lemma 5.3 implies that there is a unique Heegner divisor of type 2 and index 1. $\square$

We denote this divisor with $\mathcal{D}_{con}$. By Corollary 4.8 the Picard lattice of the generic $K3$ surface X in $\mathcal{D}_{con}$ is given by:

$$Pic(X) = \langle L_+, R_1, R'_1 \rangle.$$

The intersection matrix of the lattice $Pic(X)$ with respect to the basis $\tilde{e}_0, \tilde{e}_1, \dots, \tilde{e}_7, R_1, R'_1$:

$$Q(X) = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 2 & 4 \\ 0 & -2 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & -2 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & -2 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & -2 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & -2 & 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 & -2 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -2 & 0 & 2 \\ 2 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & -2 & 2 \\ 4 & 1 & 1 & 1 & 1 & 2 & 2 & 2 & 2 & -2 \end{pmatrix}$$

5.3. Quartics with splitting cubics. A simple remark is the following:

Lemma 5.5. *A plane quartic C has a hyperflex line if and only if there exists a cubic D such that the 4:1 cover of $\mathbb{P}^2$ branched along C is trivial over D .*

Proof. Notice that a quartic C has a hyperflex line L if and only if there is a point $p \in C$ with

$$4p \in |O_C(1)| = |K_C|.$$

In this case, let M be a line through p , $M \neq L$, then $M \cdot C - p = p_1 + p_2 + p_3$. Hence

$$4(p_1 + p_2 + p_3) \in |4K_C - 4p| = |3K_C|$$

i.e. there exists a cubic curve D intersecting C with multiplicity 4 in each point and such that the points in $C \cdot D$ lie on the line M . Equivalently, by Proposition 5.1, the 4:1 cover of $\mathbb{P}^2$ branched along C is trivial over D .

Conversely, if there exists a cubic D with $D \cdot C = 4(p_1 + p_2 + p_3)$ and a line M with $M \cdot D = p_1 + p_2 + p_3$ then

$$M \cdot C = p_1 + p_2 + p_3 + p,$$

hence $4p = 4M \cdot C - D \cdot C \in |O_C(1)|$ i.e. p is a hyperflex point. $\square$

Remark 5.6. In fact, Lemma 5.5 follows immediately if we recall that, up to a projective transformation, the equation of a quartic C with a hyperflex line is of the form:

$$xf_3(x, y, z) + z^4 = 0,$$

where f_3 is a cubic polynomial in $\mathbb{C}[x, y, z]$.

In general, a (singular) plane cubic D can be a splitting curve for a quartic C even if the 4:1 cover branched along C is not trivial over D i.e. the irreducible components of $\pi^{-1}(D)$ are not isomorphic to D . Let $\pi : X_C \rightarrow \mathbb{P}^2$ and $\pi_1 : S_C \rightarrow \mathbb{P}^2$ be the 4:1 and the double cover branched along C respectively.

Proposition 5.7. *Let C be a plane quartic with a splitting nodal cubic D . Then X_C belongs to a Heegner divisor of index 2 and*

- i) type 3 if π is trivial over D (i.e. X_C lies in $\mathcal{D}_{flex}$),*
- ii) type 7 if π_1 is trivial over D ,*
- iii) type 5 otherwise.*

Proof. Up to projectivities we can assume that D is defined by the equation:

$$y^2 = x^3 + x^2.$$

A parametrization for D is given by:

$$\phi : \mathbb{C} \longrightarrow D, \quad t \longmapsto (t^2 - 1, t(t^2 - 1)),$$

where $\phi(1) = \phi(-1)$ is the node of D .

The plane quartic C with equation $f_4(x, y, z) = 0$ intersects D in the points $\phi(t)$, with:

$$f_4(t^2 - 1, t(t^2 - 1), 1) = 0.$$

In particular, if the intersection multiplicity of each point in $C \cap D$ is a multiple of four, then:

$$f_4(t^2 - 1, t(t^2 - 1), 1) = q^4,$$

where $q \in \mathbb{C}[t]$ and $q^4 \in \mathbb{C}[t^2 - 1, t(t^2 - 1)]$. It can be easily proved that

$$\mathbb{C}[t^2 - 1, t(t^2 - 1)] = \{p \in \mathbb{C}[t] : p(1) = p(-1)\}.$$

Hence:

$$q(1) = i^a q(-1), \quad a = 0, \dots, 3.$$

The 4:1 cover of the plane branched along C is given by:

$$X_C = \{w^4 = F(x, y, z)\} \subset \mathbb{P}^3.$$

Notice that the map ϕ lifts to four distinct maps to X_C :

$$\phi_k : \mathbb{C} \longrightarrow X_C, \quad t \longmapsto (t^2 - 1 : t(t^2 - 1) : 1 : i^k q(t)),$$

hence the inverse image of D in X_C has four isomorphic irreducible components:

$$D_k = \phi_k(\mathbb{C}), \quad a = 0, \dots, 3.$$

We now study the restriction of the cover π to D for all values of a .

i) If $a = 0$, then $q(1) = q(-1)$, hence each component D_k is a nodal curve and π is trivial over D . By Lemma 5.5 the plane quartic C has a hyperflex line and X_C belongs to the Heegner divisor $\mathcal{D}_{flex}$.

ii) If $a = 2$, then $q(1) = -q(-1)$, hence the curves D_k are smooth rational curves. Note that $\phi_0(1) = \phi_2(-1)$ and $\phi_0(-1) = \phi_2(1)$, so D_0 and D_2 meet in two points over the singular point $p = (0 : 0 : 1) \in D$. Similarly, D_1 and D_3 meet in two points over p . In this case the double cover π_1 of $\mathbb{P}^2$ branched along C is trivial over D , in particular the inverse image of D in S_C is the union of two singular curves. Note that:

$$r = D_0 - \tau^*(D_0) = D_0 - D_2 \in \text{Pic}(X_C) \cap L_-$$

and $r^2 = -4 - 2 \cdot 2 - 2 \cdot 3 = -14$. We have a lattice embedding:

$$\Lambda = \langle r, \sigma^*(r) \rangle \subset \text{Pic}(X_C) \cap L_- \cong A_1(n)^{\oplus 2}.$$

In particular $-14 = -2n(a^2 + b^2)$, $a, b \in \mathbb{Z}$. Thus we have $n = 7$ and X_C belongs to a Heegner divisor of type 7 and index 2.

iii) If $a = 1$, then $q(1) = iq(-1)$, hence the curves D_k are smooth. In this case we have also $\phi_0(1) = \phi_1(-1)$ and $\phi_0(-1) = \phi_3(1)$, so D_0 meets D_1 and D_3 . Similarly, D_2 meets D_3 and D_1 . In particular, the inverse image of D in S_C is the union of two smooth rational curves. Note that:

$$r = D_0 - \tau^*(D_0) = D_0 - D_2 \in \text{Pic}(X_C) \cap L_-$$

and

$$r^2 = -4 - 2 \cdot 3 = -10.$$

Then:

$$\Lambda = \langle r, \sigma^*(r) \rangle \subset \text{Pic}(X_C) \cap L_-.$$

In fact (since $-r^2/2$ is prime and $n \neq 1$) we easily get that this is an equality. Hence X_C belongs to a Heegner divisor of type 5 and index 2. This could be proved also by applying Proposition 4.9 (note that the degree is minimal since $m = 2$ and we can assume that C has no hyperflex lines).

iv) The case $a = 3$ is analogous to case iii). $\square$

Remark 5.8. A similar description could be given more generally for any rational splitting curve.

APPENDIX A. LATTICES

A *lattice* L is a free $\mathbb{Z}$ -module of finite rank endowed with an integral symmetric bilinear form

$$(\cdot, \cdot) : L \times L \longrightarrow \mathbb{Z}.$$

Let $r(L)$ be its *rank*, $d(L)$ its *discriminant* and $s(L) = (s_+, s_-)$ its *signature*. We denote with $O(L)$ the group of self-isometries of L .

If L_1, L_2 are two lattices then $L_1 \oplus L_2$ is their *orthogonal direct sum* and $L^{\oplus n}$ is the orthogonal direct sum of n copies of L . Let a be an integer, then $L(a)$ is the lattice with the same underlying $\mathbb{Z}$ -module of L and with bilinear form multiplied by a .

In the following, we assume that L is an even lattice i.e. (x, x) is even for each $x \in L$. Let $L^* = \text{Hom}(L, \mathbb{Z})$ be the dual lattice of L , then there is a canonical embedding $L \subset L^*$. The *discriminant group* of L is the quotient group

$$A_L := L^*/L.$$

The group A_L is an abelian group of order $|d(L)|$, we denote by $\ell(L)$ the minimal number of its generators. The bilinear form on L can be extended to L^* with values in $\mathbb{Q}$ and the *discriminant quadratic form* of L is the map defined by

$$q_L : A_L \longrightarrow \mathbb{Q}/2\mathbb{Z}, \quad q_L(x + L) = (x, x) + 2\mathbb{Z}.$$

Examples:

- i) The *hyperbolic plane* is the rank two even lattice $U = \langle x, y \rangle$ with bilinear form: $(x, x) = 0, (x, y) = 1$. This is an even unimodular lattice of signature $(1, 1)$.
- ii) *Lattices of type ADE*. We denote by A_m, D_n, E_l , ($m \geq 1, n \geq 4, l = 6, 7, 8$) the negative definite even lattices associated with the corresponding Cartan matrixes. By the classification theorem of root systems (see [3]), any negative definite lattice generated by elements with square -2 is an orthogonal sum of lattices of type *ADE*. The discriminant groups of these lattices are:

Proposition A.1 (Proposition 3.2.2, [10]).

$$\begin{aligned} A_{A_m} &\cong \mathbb{Z}/(m+1)\mathbb{Z}; \\ A_{D_n} &\cong \begin{cases} \mathbb{Z}/4\mathbb{Z}, & \text{if } n \text{ is odd,} \\ (\mathbb{Z}/2\mathbb{Z})^2, & \text{if } n \text{ is even;} \end{cases} \\ A_{E_l} &\cong \mathbb{Z}/(9-l)\mathbb{Z}; \end{aligned}$$

where $m \geq 1, n \geq 4, l = 6, 7, 8$.

An even lattice S is *2-elementary* if $A_S \cong (\mathbb{Z}/2\mathbb{Z})^{\alpha_S}$ for some non negative integer α_S . We can associate to S the following invariant:

$$\delta_S = \begin{cases} 0 & \text{if } q_S(x) \in \mathbb{Z} \text{ for all } x \in A_S \\ 1 & \text{otherwise} \end{cases}$$

The following result gives an answer to the problem of classification of 2-elementary lattices:

Theorem A.2 (Theorem 3.6.2, [11]). *The isomorphism class of an even 2-elementary lattice S is determined by the invariants $(s_+, s_-, \alpha_S, \delta_S)$ if $s_+ > 0$ and $s_- > 0$.*

Examples of 2-elementary lattices are given by unimodular lattices, “twisted” unimodular lattices (e.g. $U(2)$, $E_8(2)$), A_1 and D_{2n} , $n \geq 2$. We describe in detail two cases:

i) Let $A_1 = \langle x \rangle$, then $A_{A_1} = \langle x/2 \rangle \cong \mathbb{Z}_2$ and $q_{A_1}(x/2) = -1/2$. In particular $\alpha_{A_1} = \delta_{A_1} = 1$. This lattice has a (unique up to isometries) structure of free $\mathbb{Z}[i]$ module given by the action of the isometry:

$$J_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

ii) The lattice D_4 can be described as:

$$D_4 = \langle \pm e_i \pm e_j \rangle, \quad i, j = 1, \dots, 4,$$

where $(e_i \cdot e_j) = -\delta_{ij}$. In fact, a basis is given by:

$$f_1 = e_1 - e_2, \quad f_2 = e_3 - e_4, \quad f_3 = e_2 - e_3, \quad f_4 = e_3 + e_4.$$

We have

$$A_{D_4} = \langle e_1, (e_1 + e_2 + e_3 + e_4)/2 \rangle = \langle (f_2 + f_4)/2, (f_1 + f_2)/2 \rangle$$

(where we write representatives of the classes). The quadratic form, with respect to the given basis for A_{D_4} is given by:

$$\begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}.$$

In particular $\alpha_{D_4} = 2$ and $\delta_{D_4} = 0$. The lattice D_4 is a free $\mathbb{Z}[i]$ -module of rank two given by the action of the isometry:

$$J_2 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 \\ -1 & 1 & 2 & 1 \end{pmatrix}$$

with respect to the basis $f_1, \dots, f_4$ (it can be proved that this structure is unique up to isometries). More easily, the action on e_1, e_3, e_2, e_4 is given by the matrix $J_1 \oplus J_1$.

Remark A.3. Let $r = a_1 e_1 + \dots + a_4 e_4$, $\sum a_i \equiv 0 \pmod{2}$ be a primitive vector in D_4 . If r is primitive in D_4 (in particular not all a_i 's are even) we have the cases:

- i) If a_i is odd for $i = 1, \dots, 4$ then $r/2$ is non-trivial in A_{D_4} .
- ii) If a_1, a_3 are odd and a_2, a_4 are even, then $(r + \sigma^*(r))/2$ has integer coefficients but either 1 or 3 of the coefficients are odd, hence it is a non-trivial element in A_{D_4} .
- iii) If a_1, a_2 are odd and a_3, a_4 are even, then all coefficients of $(r + \sigma^*(r))$ are odd integers, so $(r + \sigma^*(r))/2$ is non-trivial in A_{D_4} .

An embedding of lattices $S \subset L$ is *primitive* if L/S is a free $\mathbb{Z}$ -module. Two primitive embeddings $S \subset L$ and $S \subset L'$ are *isomorphic* if there exists an isomorphism between L and L' which is the identity on S .

Let $S \subset L$ be a primitive embedding of even lattices and $K = S_L^\perp$ be the orthogonal lattice to S in L . Then $S \oplus K \subset L$ and

$$H_L = L/(S \oplus K) \subset A_S \oplus A_K$$

is isotropic. Moreover:

$$q_L = (q_S \oplus q_K) \mid H_L^\perp/H_L.$$

Let p_S, p_K be respectively the projections of H_L to A_S and A_K . By the primitivity of the embedding, these projections are embeddings:

$$p_S : H_L \longrightarrow H_L(S) \subset A_S, \quad p_K : H_L \longrightarrow H_L(K) \subset A_K.$$

The group H_L is the graph of the isomorphism

$$\gamma_{S,K}^L = p_K \circ p_S^{-1} : H_L(S) \longrightarrow H_L(K)$$

in $A_S \oplus A_K$. Since H_L is isotropic we also have:

$$q_K \circ \gamma_{S,K}^L = -q_S.$$

This construction can be also reversed.

Proposition A.4 (Proposition 1.6.1, [10]). *A primitive embedding of an even lattice S into another even lattice L with discriminant form q_L and orthogonal complement isomorphic to K is determined by a pair (H, γ) , where:*

- i) $H \subset A_S$ and $\gamma : H \longrightarrow A_K$ is a group monomorphism,
- ii) $q_K \circ \gamma = -q_S \mid H$,
- iii) $q_L \cong (q_S \oplus q_K \mid \Gamma^\perp/\Gamma)$ where Γ is the graph of γ in $A_S \oplus A_K$.

Two such pairs (H, γ) and (H', γ') determine isomorphic primitive embeddings if and only if $H = H'$ and γ, γ' are conjugate via a self-isometry of K .

In particular, if L is unimodular, the embedding is determined by an isomorphism $\gamma : A_S \longrightarrow A_K$ satisfying ii).

REFERENCES

- [1] M. Artebani. A compactification of $\mathcal{M}_3$ via $K3$ surfaces. *preprint*, 2005.
- [2] W. Barth, C. Peters, and A. Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 1984.
- [3] N. Bourbaki. *Éléments de mathématique. Fasc. XXXIV. Groupes et algèbres de Lie. Chapitre IV: Groupes de Coxeter et systèmes de Tits. Chapitre V: Groupes engendrés par des réflexions. Chapitre VI: systèmes de racines*. Hermann, Paris, 1968.
- [4] I. Dolgachev and D. Ortland. Point sets in projective spaces and theta functions. *Astérisque*, (165):210 pp., 1988.
- [5] I. Hadan. Tangent conics at quartic surfaces and conics in quartic double solids. *Math. Nachr.*, 210:127–162, 2000.
- [6] S. Kondō. The moduli space of 8 points on $\mathbb{P}^1$ and automorphic forms. *math.AG/0504233*.
- [7] S. Kondō. A complex hyperbolic structure for the moduli space of curves of genus three. *J. Reine Angew. Math.*, 525:219–232, 2000.
- [8] E. Looijenga. Compactifications defined by arrangements. I. The ball quotient case. *Duke Math. J.*, 118(1):151–187, 2003.
- [9] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric Invariant Theory*. Ergeb. Math. Grenzgeb. (2), 34, Springer, 1994.
- [10] V. V. Nikulin. Factor groups of groups of the automorphisms of hyperbolic forms with respect to subgroups generated by 2-reflections. *J. Soviet Math.*, 22:1401–1475, 1983.

- [11] V.V. Nikulin. Integral symmetric bilinear forms and its applications. *Math. USSR Izv.*, 14:103–167, 1980.
- [12] A.M. Vermeulen. *Weierstrass points of weight two on curves of genus three*. PhD thesis, Universiteit van Amsterdam, 1983.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI MILANO, VIA C. SALDINI 50, MILANO, ITALIA
E-mail address: `michela.artebani@unimi.it`